

maxima per replicate were on the order of 350-450 flies.

Every other generation 100 female flies from each replicate were assayed for sternopleural bristle count. Samples from males invariably averaged a few percent lower in bristle count than females. The 22nd generation results, based on the pooled counts of 600 females taken from subreplicates of replicates, are summarized below.

Means and 95% Confidence Intervals

Line I Replicate		Line II		Line III	
1	17.43±.22	1	17.83±.22	1	16.79±.21
2	17.65±.25	2	17.72±.23	2	16.58±.16
3	17.64±.23	3	17.96±.23	3	18.58±.26

Inter- and intra- line divergence are relatively slight in Lines I and II. The initiation of Line II populations with small samples of the parental gene pool did not lead to a drifting apart of bristle count. The more drastic decimation regimen of Line III was effective in producing drift. Two generation 22 replicates have counts significantly less than any Line I or II replicates, and one replicate is significantly higher. A plot of replicate bristle count against generations (not shown) indicates that the dispersal of Line III replicates developed gradually and was still increasing at the termination of the experiment.

Monclús, M. University of Barcelona, Spain. Influence of day time and season on mating propensity in *D. subobscura*.

A strong influence of day time and season has been detected in the mating propensity of *D. subobscura*. This relation came out in tests carried on with a different purpose. In each test 50 ♂♂ and 25 virgin ♀♀

were put together and the number of matings accomplished during one hour was recorded. Flies of different ages were tested separately, but in the results here presented all the ages are lumped.

The individuals were developed in our standard conditions of culture for *D. subobscura*, in a room with controlled temperature at $17^{\circ} \pm 0.5^{\circ}$ C. The mating tests were performed in all seasons at 22° or 23° C. The stock used has been kept in the laboratory for two years.

Routine tests were performed at 11 a.m. since December to June. Working at the same time of the day in July and August it became difficult to get results because of the very few matings observed. Since *D. subobscura* in the natural populations is active in summer only early in the morning and in the evening, the time of testing was moved to 6:45 a.m. The mean mating frequencies observed in the tests carried out in these three different conditions, are as follows:

December-June	11	a.m. (32 tests)	M = 16.03	matings for test		
July-August	11	a.m. (13 tests)	M = 1.30	"	"	"
July-August	6:45	a.m. (9 tests)	M = 10.44	"	"	"

These results seem to indicate that the sexual activity of *D. subobscura* is controlled by an internal rhythm, perhaps related to some external factor difficult to identify.

Robertson, F. W. and Chipchase, M. Department of Genetics, University of Edinburgh. The comparison of genetic differences by hybridization between DNA and RNA synthesized in vitro.

DNA prepared from different species of *Drosophila* has been used as template to synthesize complementary RNA (c-RNA) by RNA polymerase extracted from *Micrococcus lysodeikticus*. The general properties of the hybridization between such DNA and RNA have been studied and the RNA transcribed

from *melanogaster* template has been annealed with DNA from various species to determine the level of discrimination. The ribonuclease resistant RNA, bound to denatured DNA, is recovered on membrane filters and separate labelling of the DNA and RNA has been used to estimate the fraction of the DNA which is bound to RNA. The level of hybridization between *D. melanogaster*