

Lamborot, M. and S. Koref-Santibañez.  
Universidad de Chile, Santiago, Chile.  
Temperature and sexual isolation  
between *D. gaucha* and *D. pavani*.

A study of mating activity of the sibling  
species *D. pavani* and *D. gaucha* (DIS 1966 42:  
106; *Biológica* 1967 61: 3-6) revealed that the  
optimum temperature for sexual activity was  
lower (18°C) in the Chilean than in the Brazil-  
ian species (25°C). Interspecific crosses

showed a differential receptivity of the females. As the isolation indices could not be obtained, a new set of experiments was set up, using the "male choice method" at nine different temperatures: 6°C, 8°C, 12°C, 16°C, 20°C, 24°C, 28°C, 32°C and 34°C. For each temperature the activity of approximately 100 males was studied. 5 males were placed for 6 hours with 5 females of their own and 5 of the sibling species in 5 x 20 cm. vials. The ventral receptacle and the spermathecae of the females were examined for the presence of sperm.

Tables 1 and 2 summarize the results obtained, together with the isolation coefficients

Table 1. ♂ *pavani*

| Temp. °C | Homog. % | Heterog. % | K         |
|----------|----------|------------|-----------|
| 6        | 32.0     | 16.0       | 0.43±0.19 |
| 8        | 48.9     | 34.8       | 0.22±0.11 |
| 12       | 42.3     | 39.6       | 0.05±0.08 |
| 16       | 71.2     | 72.5       | 0.01±0.07 |
| 20       | 77.3     | 79.2       | 0.01±0.07 |
| 24       | 72.5     | 83.2       | 0.16±0.08 |
| 28       | 59.6     | 61.5       | 0.02±0.09 |
| 32       | 23.6     | 34.6       | 0.22±0.04 |
| 34       | 6.0      | 24.0       | 0.63±0.01 |

Table 2. ♂ *gaucha*

| Temp. °C | Homog. % | Heterog. % | K         |
|----------|----------|------------|-----------|
| 6        | 24.0     | 17.0       | 0.16±0.22 |
| 8        | 48.9     | 17.2       | 0.51±0.10 |
| 12       | 51.2     | 21.2       | 0.49±0.07 |
| 16       | 66.2     | 22.8       | 0.61±0.06 |
| 20       | 75.7     | 27.9       | 0.59±0.06 |
| 24       | 89.1     | 72.2       | 0.27±0.08 |
| 28       | 79.4     | 40.4       | 0.45±0.08 |
| 32       | 71.8     | 20.0       | 0.69±0.07 |
| 34       | 40.4     | 4.7        | 0.85±0.14 |

Homogamic, heterogamic preferences and isolation index (K) of *D. gaucha*  
and *D. pavani* males at different temperatures

(K) (Malogolowkin-Cohen et al. *Evolution* 19: 95-103). While *D. pavani* males show little isolation throughout the temperature range, having even greater preferences for the foreign female, *D. gaucha* males reveal marked preferences for their own females, specially notorious at extreme temperatures; at the temperature at which the activity of *D. gaucha* was optimal, isolation was lowest. The results point to the conclusion that female receptivity seems to be more responsible for sexual isolation than the activity of the male.

Kuroda, Y. National Institute of Genetics,  
Misima, Japan. Effects of X-irradiation on  
the differentiation of eye-antennal discs  
of *D. melanogaster* in organ culture.

Eye-antennal discs were irradiated with 0 R,  
500 R, 1,000 R, 1,500 R and 2,000 R of X-rays  
(180 kV, 25 mA, 1.0 mm Al filter, distance  
40 cm, dose rate 300 R/min) immediately after  
their preparation in hanging drop cultures  
from mature third-instar larvae of the Oregon-

R strain of *D. melanogaster*. After irradiation the discs were cultured in a chemically defined medium containing 10<sup>-4</sup> mg/ml rubrosterone and examined for the effects of X-rays on the differentiation of ommatidia. When eye-antennal discs were irradiated with 500 R or 1,000 R no marked inhibition was observed in the differentiation of ommatidia after 24 hours of cultivation. The organization of ommatidium-forming cells into cell clusters was observed in the eye disc portion as seen in eye-antennal discs in non-irradiated control cultures. With 1,500 R the differentiation of ommatidia was partially inhibited 24 hours after explantation. 2,000 R inhibited almost completely the differentiation of ommatidia when examined after 24 hours of cultivation.

When eye-antennal discs were irradiated first with a dose of 1,000 R immediately after explantation, then they were exposed to a second dose of 1,000 R at 2 or 4 hours after explantation, the effects of X-ray were found to be different depending on the extent of the intervals between the first and second doses. With the second dose given at 2 hours after explantation the differentiation of ommatidia was partially inhibited after 24 hours of cultivation; whereas with the second dose given at 4 hours after explantation no inhibitory effect of X-ray was observed on the differentiation of ommatidia.