
Wong, P.T. and W.E. Trout III. City of Hope National Medical Center, Duarte, California. The psychic fly fish feeder: A reply to Ogden Nash.




"The Lord in His wisdom made the fly And then forgot to tell us why." The Fly, by Ogden Nash

A culture of the neurological mutant Hk¹ (who falls over when he sees something moving), is placed in container (A), in bottomless bottle (B), plugged with cotton (C), which is suspended at the surface of the water in aquarium (D). When a fly emerges (E) and sees fish (F) moving below him, he loses control of himself and falls into the water (G) thereby feeding fish (F).

Sparrow, J.C. University of Sussex, Brighton, England. Eggwashing apparatus.

This apparatus replaces the reversing pump described by Sang (1956) in our Drosophila egg sterilization procedure for axenic culture. The eggs are col-

lected by the method of Sang (1956), dechorionated with hypochlorite and suspended in 1.0% saline. This stops the dechorionated eggs from aggregating. Any larvae are removed by hand using a binocular microscope. The eggs are then placed in the apparatus through the thistle funnel and enter the egg chamber. They are washed in this chamber by a sterile 0.1% C.D.B.A.C. (cetyldimethylbenzylammonium chloride) solution which runs continuously through the apparatus. The flow rate is controlled by the screw clip (S) at a rate of 90 drops/minute



Egg Chamber