

Ogah, F. and R. MacIntyre. Cornell University, Ithaca, New York. The persistence of Acid phosphatase-1 null mutants of *Drosophila melanogaster* in experimental population cages.

Four Bennett box type population cages were set up to study the fate of Acph-1^n null alleles in laboratory populations. Two null allele stocks Acph-1^{n-5} and Acph-1^{n-11} (Bell and MacIntyre, 1972) obtained by means of EMS (Bell and MacIntyre, 1972) were used along with wild type stocks containing the fast allele, Acph-1^B , to

set up the cage populations as shown in Table 1.

The initial frequencies of Acph-1^B are 0.80 in cages 1 and 2, and 0.20 in 3 and 4. F_1 heterozygotes as founders were used to minimize the initial effect of "luxuriant" heterosis

Table 1. Founders of the cages.

Cages 1 and 2		Cages 3 and 4	
120 ♂♂		60 ♂♂	
120 ♀♀ virgins	BB stock (wild type)	60 ♀♀ virgins	null stock (n-5)
40 ♂♂ F_1	null stock (n-5) x	60 ♂♂	
40 ♀♀ virgins F_1	BB (wild type)	60 ♀♀ virgins	null stock (n-11)
40 ♂♂ F_1	null stock (n-11) x	40 ♂♂ F_1	null stock (n-5) x
40 ♀♀ virgins F_1	BB (wild type)	40 ♀♀ virgins F_1	BB (wild type)
		40 ♂♂ F_1	null stock (n-11) x
		40 ♀♀ virgins F_1	BB (wild type)

in cages founded by flies from different strains. (See MacIntyre and Wright, 1966).

For the first eleven months egg samples were taken monthly from each cage. A thin layer of food was spread on the handle of plastic teaspoons, and one of these was put in a vial and placed into the cage in the normal position of a food vial. This was allowed to stay for thirty six hours. Then the thin layer of food was transferred to a half pint bottle where the eggs were cultured. The adult flies were collected and acid phosphatase activity was assayed for by means of spot test (Bell, et al., 1972). In this assay the null homozygotes are completely lacking the bright red color indicative of Acph-1^B homozygotes and $\text{Acph-1}^n/\text{Acph-1}^B$ heterozygotes. Consequently, only null homozygotes could be used to determine the Acph-1^n frequencies in the cages.

Table 2 shows the trend in the change of the frequencies of null alleles from one generation to the next for the four population cages. Initially in cages 1 and 2 the frequencies of null mutant rose slightly, and those for 3 and 4 decreased slightly. Then there was an apparent stabilization at frequencies of about 0.6 for null alleles in all the cages.

Table 2. Frequency of Acph-1^n

Months	Date	CAGES			
		1	2	3	4
1	7/26/70	0.36 ± 0.03	---	0.74 ± 0.03	---
2	8/26/70	0.46 ± 0.05	0.36 ± 0.03	0.67 ± 0.03	0.78 ± 0.02
3	9/26/70	0.14 ± 0.03	0.12 ± 0.02	0.58 ± 0.05	0.75 ± 0.04
4	10/26/70	0.39 ± 0.03	0.18 ± 0.03	0.29 ± 0.03	0.73 ± 0.02
5	11/27/70	0.65 ± 0.04	0.41 ± 0.04	0.69 ± 0.04	0.66 ± 0.03
6	12/26/70	0.49 ± 0.04	0.55 ± 0.04	0.66 ± 0.04	0.68 ± 0.04
7	1/26/71	0.41 ± 0.04	0.60 ± 0.06	0.76 ± 0.03	0.57 ± 0.03
8	2/26/71	0.37 ± 0.04	0.48 ± 0.04	0.59 ± 0.03	0.62 ± 0.04
9	3/26/71	0.42 ± 0.04	0.57 ± 0.04	0.69 ± 0.03	0.61 ± 0.04
10	4/26/71	0.50 ± 0.04	0.45 ± 0.04	0.50 ± 0.05	0.62 ± 0.04
11	5/26/71	0.44 ± 0.04	0.35 ± 0.04	0.41 ± 0.04	0.52 ± 0.04
24	6/12/72	0.46 ± 0.05	0.54 ± 0.05	0.49 ± 0.05	0.35 ± 0.04

The apparent lack of selection against Acph-1^n homozygotes and heterozygotes in experi-

Berger, E.M. and R. Canter. State University of New York, Albany, N.Y.
"Moulting" esterases of *Drosophila*.

6 through 10 are stage specific. Esterases 8, 9 and 10 are present in first instar larvae, but gradually diminish in activity, so that

During the ontogeny of *Drosophila pseudoobscura*, a series of 10 β -naphthyl esterase isozymes can be observed by electrophoresis in acrylamide gels. As seen in Figure 1, esterases 1 through 5 appear throughout development, while esterases 6 through 10 are stage specific. Esterases 8, 9 and 10 are present in first instar larvae, but gradually diminish in activity, so that by the third instar they are absent. At the onset of pupation these isozymes reappear, and continually increase in activity through metamorphosis. Esterase 6 and 7 first appear during adult devel-




Fig. 1. Ontogeny of esterase isozymes in *D. pseudoobscura*. Stages are fli - first larval instar; sli - second larval instar; tli - third larval instar; whp - white pupa; tnp - tan pupa, no eye color; rep - pupa, eye colored red, no other pigmentation; pep - pre-emergent pupa, fully pigmented; ad - 9 day old adult; pc - discarded pupal case. *D. pseudoobscura* strain Est 51.12 used in this study was obtained from Dr. R.C. Lewontin, Dept. of Biology, University of Chicago.

opment, increasing in activity until eclosion. At eclosion esterases 6, 7, 8, 9 and 10 simultaneously disappear from the adult. The loss of these isozymic forms is not due to some rapid inactivation, for all five isozymes can be observed in homogenates of the discarded pupal case.

Esterases 1 through 5 are absent in these homogenates. Cohomogenization of newly eclosed adults and their empty pupal case restores the 10 banded pre-eclosion pattern.

While the function of these stage specific esterases is unknown, their developmental timing and post-eclosion localization, suggest a possible role in the moulting process: we therefore designate these enzymes as "moulting esterases". Analogous esterase systems have now been observed in *D. melanogaster*, *D. robusta*, *D. pattersoni*, *D. buskii* and *D. funebris*. In *D. pseudoobscura* we now have allelic variants for several of these systems and are proceeding in genetic analyses.

This work has been supported by a grant from the NIH (GM 18910).

(Continued from preceding page)

mental populations is rather surprising. In addition, other tests indicate that the viability and developmental rate of null mutants do not differ from those of wild type flies. The following question, then, might be asked: Is acid phosphatase-1 a relict enzyme? No naturally occurring null alleles have been found in populations of *D. melanogaster* and, in fact, in this species, the gene-enzyme system is very monomorphic (O'Brien and MacIntyre, 1969; MacIntyre, 1972 and unpublished). The observation that approximately 90% of the soluble acid phosphatase activity detected by our assay methods is due to acid phosphatase-1 also argues against a trivial role for this enzyme in *D. melanogaster*. (Bell and MacIntyre, 1973.)

The apparent dilemma may be due, in part, to the undoubtedly real differences between the environments of the laboratory and the fly's natural ecological niche. Certain factors which make this enzyme activity very important may be absent under laboratory conditions. We plan to test this possibility as soon as the physiological substrate(s) of the enzyme are determined.

The results described here have another important ramification. If the environment of the laboratory is so different that selection pressure against individuals deficient for a particular enzyme is absent or even reduced, experiments designed to detect selective differences between individuals carrying different allozymic forms of that enzyme, are almost certainly going to give negative results. (For example, see Yamazaki, 1971.) In other words, a demonstration of selective neutrality of certain alleles specifying allozymes is, in fact, a meaningless result if the organism doesn't even need the enzyme under the conditions of the experiment.