

Table 1 summarizes some of our experiences with this technique, showing the averaged results from three separate samples, each of which involved a 1 hr. pre-collection followed by an egg collection of the indicated time. It is evident that the production of eggs peaks at 1 hr. and appears to slowly decline thereafter. If the pre-collection has been sufficient, then the percentage of synchronously developing embryos stays very high and only decreases after 2 hours. The percentage of abnormally developing embryos also increases as collection time increases but never exceeds 10%. If synchronously developing embryos are not an important consideration, one can use longer egg laying periods (Table 2) and obtain much larger

Table 2.

Hours	Collection	No. females per bottle	No. eggs per plate		No. eggs per female per hr.
			Range	Mean	
0-2	Pre-collection	120	2000-4000	3000	13
2-4	Collection #1	120	1000-2000	1500	7
4-6	Collection #2	120	500-1000	750	4

numbers of eggs per plate. It can be seen, however, that egg production per female decreases linearly to very low levels by 6 hours. There-

fore, females are not pushed for more than 4-6 hours of continuous collections. In our hands, then, the optimal collection time is one hour, which will produce a high level of synchrony, minimal numbers of abnormal embryos and a moderately large number of eggs per plate.

In a 4 hour day it is, theoretically, possible to collect 20,000-40,000 eggs excluding the pre-collection. In practice, this technique has been able to produce a minimum of 28,000 embryos in an 8-day period. This figure is a minimum estimate since it does not include an approximately equal number of eggs which were discarded as surplus. These collections were done using anywhere from 8-15 collection bottles which held the adults obtained from 15-30 food bottles. This gave us an average yield of about 3,500 eggs per day, a figure which is in general agreement with the data contained in Table 1. As a Drosophilist may know, egg production is variable and there have been times when we have obtained little or no eggs; conversely, there have been times when we have obtained 40,000 eggs in one collection. In summary, we have devised a technique which is as productive as a population cage, and which can produce a moderately large number of eggs using a smaller number of adults, and which is more responsive to varying daily needs for embryos.

Pasic, T.R. and H. Nickla. Creighton University, Omaha, Nebraska. A BASIC program for construction of fate maps.

A computer program in BASIC has been developed to aid in the construction of blastula "fate maps". Early mitotic misdivisions in the embryo can be induced by an unstable ring X-chromosome. This results in two clones of nuclei ran-

domly oriented with respect to the blastula cortex and differing in X-chromosome complement. When the nuclei migrate and become fixed to the cortex, a random plane of division divides the blastula. The probability of a plane of division falling between any two imaginal discs increases as the distance between two imaginal discs increases (Hotta and Benzer 1972).

Adult mosaics can be scored for any number of surface landmarks on the basis of tissue type (1 = male, 2 = female, 3 = mixed). Each scoring is entered into a data table at the location (A,B) where A is a unique integer for each fly (1,2,3, . . .) and B represents each landmark in coded form (1 = eye, 2 = wing, etc.). The computer effects a two-by-two pairing of landmarks and counts the number of times that the two scorings are different. This sum represents the number of times a plane of division fell between the imaginal discs. The sum is converted into a percentage and prints it as the distance (in Sturts) between each pair of imaginal discs. (Note: The total number of pairings is $(N^2 - N)/2$ where N is the total number of landmarks.)

Other subroutines in the program include the distance between homologous landmarks; total percent male tissue, percent male tissue at each landmark and percent male tissue of each fly. The program was tested using data on 196 mosaics having 39 surface landmarks. The program is available upon request.

Reference: Hotta, Y. and S. Benzer 1972, Nature 240:527.