

sperm. Manipulation of the shape of the productivity curve through manipulation of culturing conditions should make it possible to test some predictions of the nature of sperm competition, as detailed elsewhere (Gromko in prep.).

References: Gilbert, D.G., R.C. Richmond & K.B. Sheehan 1981, *Evolution* 35:21-37; Gromko, M.H. in prep., A new model of sperm competition in *Drosophila melanogaster*; Pyle, D.W. & M.H. Gromko 1978, *Experientia* 34:449-450.

Guest, W.C. University of Arkansas, Fayetteville. Chlorpromazine delays *D. melanogaster* larval development.

Chlorpromazine (CPZ) is a widely used tranquilizer that is thought to block dopamine neuro-receptor sites in the brain of vertebrates (Gale 1980) and may act in other ways as well. Dopamine is an intermediate in the synthesis of the tanning pigment sclerotin involved in the molting of insects (Karlson & Sekeris 1966). When *D. melanogaster* first instar larvae were fed 0.2 mg/ml CPZ in laboratory food pupation was delayed approximately three days. At a concentration of 0.3 mg/ml there was a delay in pupation of five and one-half days although at this concentration only three percent of the larvae survived to pupate.

When second instar larvae were treated with CPZ the delay in pupation increased with the concentration of CPZ. The delay varied from four days at 0.2 mg/ml to seven days at 0.1 mg/ml. There was no reduction in survival at 0.2 mg/ml but at 0.6 mg/ml only 47 percent of the larvae survived and at 1.0 mg/ml there was only a six percent survival rate. When third instar larvae were treated there was a delay in pupation of approximately three days at all concentrations up to 2.0 mg/ml and the percent survival varied directly with the concentration from 80 percent survival at 0.2 to 16 percent at 2.0 mg/ml.

There are no reports in the literature on the effects of CPZ on insect larvae. Studies with vertebrates indicates that the drug may interfere with steroid hormone function (Wakabayashi et al. 1980), block dopamine receptors (Gale 1980), as well as decrease membrane permeability (Maoi 1979). Most investigators have indicated that a block in dopamine utilization occurs and this appears to be a reasonable explanation of the action of CPZ in insect development. The availability of dopamine to form sclerotin would have an adverse effect on pupation.

References: Gale, K. 1980, *Nature* 280:576-580; Karlson, P. & C. Sekeris 1966, *Acta Endocrin.* 53:505-518; Maoi, M., T. Suzuki & K. Tagi 1979, *Biochem. Pharmacol.* 28:295-299; Wakabayashi, I., M. Kanda, N. Miki, H. Miyoshi, E. Ohmura, D. Demura & K. Shizume 1980, *Neuroendocrinology* 30:319-322.

Gupta, A.P. Instituta Biologica da UFRJ, Rio de Janeiro, Brasil. Molecular evidence for developmental stability in species crosses and backcross progeny of *D. pseudoobscura* and *D. persimilis*.

Prakash & Merritt (1972) reported that at the adult acid phosphatase-6 (AP-6) locus, two alleles determining the presence (+) or the absence (-) of the enzyme are found in *D. pseudoobscura*, but this locus is monomorphic for the absence in the adults of *D. persimilis* (Prakash 1977). AP-6 is sex linked and the + allele is dominant over the - allele. In *D. pseudoobscura*, the frequency of the + allele is 30-40% in standard arrangement, whereas this allele is absent in the sex ratio arrangement.

Even though these two species are similar in morphology, they show significant genetic differences. The F_1 males of the species cross are sterile and backcross progeny have very low viability. The sterility in F_1 males is caused by abnormal spermatogenesis. A breakdown of developmental stability in species crosses and backcrosses occurs due to unfavorable interactions of chromosomes from the two species.

The present experiment was designed to examine the level of enzyme activity at the adult acid phosphatase-6 locus in interspecific crosses and backcrosses. Two strains of *D. pseudoobscura* homozygous for + allele and two strains of *D. persimilis* homozygous for the - allele were used. Virgin females and males were collected to make F_1 's and various backcrosses. The species identity of strains was confirmed by demonstrating the sterility of both classes of F_1 hybrid males. Ten replicates each of parental, F_1 's (in both directions) and four backcross classes (only F_1 females could be used for making backcrosses) were reared concurrently at 17.5°C. Fifty individual females from each of the parental, F_1 's and various backcross