

approximately 2 days the first flies start to die and after 3-5 days 60-80% of the females have died. In contrast the "aseptic" flies show hardly any mortality during the same period. The fact that "aseptic" flies show no mortality indicates that the mortality is caused by an interaction between micro-organisms and the food. This is supported by the results of a second experiment of which the results are presented in Table 2. This table shows the mortality of flies after 72 hours of exposure to different conditions. Conditions A and B show the same result as presented in Table 1 and indicate that a high mortality on sodium octanoate is found when the flies are "non-aseptic". From a comparison between conditions A and C it is clear that sodium octanoate has to be present to induce a high mortality. The result of condition D shows that it is not the presence of the "non-aseptic" flies themselves that is responsible for the high mortality, but by something that is introduced into the vial by "non-aseptic" flies.

These results justify the conclusion that the presence of micro-organisms and sodium octanoate in the food creates a situation which is lethal for *D. melanogaster* adults. At the moment it is not clear what really causes the dying of flies. It may be that sodium octanoate is modified by a micro-organism to a toxic compound. Another possibility is that sodium octanoate enables a particular micro-organism to grow very rapidly and that or the micro-organism itself or one of its excrements is toxic to flies at a sufficiently high concentration.

References: Bijlsma, R. 1978, *Genet. Res. Camb.* 31:227-237; Sang, J.H. 1956, *J. Exp. Biol.* 33:45-71.

Lee, T.J. Chung-Ang University, Seoul, Korea. Systematic relationships among the species of *Drosophilidae* by the proteins electrophoretic analysis.

The difference patterns of aqueous soluble proteins and systematic relationships among the species of *Drosophilidae* in Korea were investigated by means of polyacrylamide gel disc-electrophoresis.

The number of protein bands appeared to be different in the 28 species, showing the difference patterns in mobility and density of staining of proteins.

Each species contained specific proteins. From 7 to 16 protein bands appeared in the 28 species, however most of the species had 10 bands.

In the intraspecies there were no different protein bands, and also the geographical difference of the protein patterns of the same species were not observed.

The average similarities among the species by the Whitney's formula in the result of the investigation as follows: in the case of the two subfamilies *Steganinae* and *Drosophilinae* in the family *Drosophilidae* appeared about 38%; in the case of the 5 genera *Mycodrosophila*, *Liodrosophila*, *Scaptomyza*, *Lordiphosa*, and *Drosophila* in the subfamily *Drosophilinae* appeared about 43%; in the case of the 5 subgenera *Dichaetophora*, *Hirtodrosophila*, *Paradrosophila*, *Sophophora*, and *Drosophila* in the genus *Drosophila* appeared about 46%; in the case of the interspecies in the subgenus *Sophophora* appeared about 69%; and the interspecies in the subgenus *Drosophila* appeared about 59%. However, the average similarities among the 4 species of *quinaria* species group of the subgenus *Drosophila* appeared about 72%.

This experiment means that the average similarities among species revealed high degree in accordance with the lower categories.

It is to be estimated that the study on the electrophoretic patterns of the whole proteins of the *Drosophila* species was valuable to determine the affinity among the species of subfamilies, genera, subgenera, and species groups as standard indices.

References: Throckmorton, L.H. 1962b, *Univ. Texas Publ.* 6205:415; Whitney, P.J., J.G. Vaughan & J.B. Mcale 1968, *J. Exp. Botany* 19:415-426.

Marengo, N.P. C.W. Post College of Long Island University, Greenvale, New York. Fibrillar disorganization in the "A" bands of "rotated" prepupal muscles of *Drosophila melanogaster*.

The mutation abdomen rotatum (ar) was discovered and named by Beliajeff (1931). The effect of this gene on development was described by the writer and Howland (1942). The ultrastructure of normal and "rotated" prepupal muscles was